

CRAPEMYRTLE (*Lagerstroemia* spp.): Cultivar Characteristics

compiled Dr. Kim D. Coder. University of Georgia School of Forest Resources. 1/2003

Crape myrtle, crapemyrtle, crepe myrtle, crepemyrtle, and crapemrytle are all common names for an exotic Asian plant group used extensively in warm temperate areas of the globe for adding color and texture to a landscape in Summer. Because of the length of time crapemyrtles have been available for sale and breeding work, hundreds of forms of the plant have been developed. This listing was compiled to assist landscape professionals and landscape managers compare a few select qualities for many of the commonly used and grown crapemyrtles.

This list is not comprehensive and does not represent availability. This list of crapemyrtle cultivars was developed to help direct professionals to cultivar-specific information on the internet and in the literature. Cultivars which are not listed here may have better values in the landscape than listed cultivars, but at the time of publication, reliable information was not available or easily accessible regarding cultivar characteristics. This list represents information present to the public, present within university information systems, and present through experience and observation of the compiler. This list was developed for educational purposes. This list may contain errors of compilation, typographical mistakes, and observational biases. Marketing information not fully representing (or suspected of not fully representing) cultivar attributes have been dismissed where possible. This list should not be used by itself for purchasing decisions.

“Disease resistance” is on a **0-5** scale with **5** being most resistant to common foliage diseases and **0** the poorest. “Form” is the height description of the cultivar when mature, and the weeping appearance (= **w**). Height definitions are **m** = miniature (<3 feet tall), **d** = dwarf (4-8 feet), **s** = standard (9-15 feet), and **t** = tall (>16 feet). “Genetics” are the species from which the cultivar is derived, which could include *Lagerstroemia indica* (= **i**), *L. fauriei* (= **f**), *L. speciosa* [*L. matthewsii*], or a combination of multiple species through hybridization (= **h**). “Features” provide for visual impact of bark characteristics (= **b**), and for the best relative cold tolerance (= **c**), remembering crapemyrtles as a group are limited in their hardiness.

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
<i>Acoma</i> (<i>Nacoma</i>)	12	white	<small>5 = best, 0 = poor</small> 5	s	h	c
<i>Alba Nana</i> (<i>Snowbaby</i>)	5	white	1	d	i	
<i>Alipurensis</i>	25	dark mauve	3	t	h	
<i>America</i>	9	ruby red	3	s	i	
<i>Apalachee</i>	15	light lavender	5	s	h	c-b
<i>Arlington</i>	18	pink	3	t	i	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = *L. indica*; f = *L. fauriei*; h = hybrid; B = great bark; C = best relative cold hardiness



The University of Georgia

THE UNIVERSITY OF GEORGIA, THE UNITED STATES DEPARTMENT OF AGRICULTURE AND COUNTIES OF THE STATE COOPERATING. THE COOPERATIVE EXTENSION SERVICE OFFERS EDUCATIONAL PROGRAMS, ASSISTANCE AND MATERIALS TO ALL PEOPLE WITHOUT REGARD TO RACE, COLOR, NATIONAL ORIGIN, AGE, SEX OR HANDICAP STATUS. A UNIT OF THE UNIVERSITY SYSTEM OF GEORGIA. AN EQUAL OPPORTUNITY / AFFIRMATIVE ACTION ORGANIZATION.

UNIVERSITY OF GEORGIA WARNELL SCHOOL OF FOREST RESOURCES PUBLICATION FOR03-2

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
<i>Basham's Party Pink</i> (<i>Basham Pink</i>) (<i>Basham's Pink</i>) (<i>Party Pink</i>)	30	<i>light pinkish-lavender</i>	<small>5 = best, 0 = poor</small> 4	<i>t</i>	<i>f</i>	
<i>Baton Rouge</i> (<i>Beverly</i>)	3	<i>bright red</i>	1	<i>m-w</i>	<i>i</i>	
<i>Bayou Marie</i>	3	<i>pink/deep red</i>	1	<i>m-w</i>	<i>i</i>	
<i>Bi-Color</i>	5	<i>red/white</i>	1	<i>d-w</i>	<i>i</i>	
<i>Biloxi</i>	35	<i>pale pink</i>	5	<i>t</i>	<i>h</i>	<i>b</i>
<i>Blizzard</i>	5	<i>white</i>	3	<i>d</i>	<i>h</i>	
<i>Blue</i>	10	<i>purple</i>	0	<i>s</i>	<i>i</i>	
<i>Bourbon Street</i> (<i>June Marie</i>)	3	<i>medium red</i>	1	<i>m-w</i>		
<i>Byers Hardy Lavender</i> (<i>Hardy Lavender</i>)	25	<i>med. lavender</i>	3	<i>t</i>	<i>i</i>	
<i>Byers Standard Red</i>	25	<i>red</i>	3	<i>t</i>	<i>i</i>	
<i>Byers Wonderful White</i> (<i>Byars Wonderful White</i>) (<i>Byers Clear White</i>) (<i>Byers White</i>) (<i>Alabama White</i>)	30	<i>clear white</i>	2	<i>t</i>	<i>i</i>	
<i>Caddo</i>	8	<i>bright pink</i>	5	<i>d</i>	<i>h</i>	
<i>Cajun Red</i>	3	<i>med. red</i>	1	<i>m</i>	<i>i</i>	
<i>Cajun Red Beverly</i>	3	<i>rosy-red</i>	1	<i>m</i>	<i>i</i>	
<i>Candycane</i>	20	<i>bi-color - pink/white</i>	3	<i>t</i>	<i>i</i>	
<i>Candystripe</i>	20	<i>bi-color - rose pink/white</i>	3	<i>t</i>	<i>i</i>	
<i>Carolina Beauty</i>	25	<i>dark red</i>	0	<i>t</i>	<i>i</i>	
<i>Catawba</i>	16	<i>violet purple</i>	3	<i>t</i>	<i>i</i>	
<i>Cedar Red</i>	4	<i>yellowish-red</i>	2	<i>d</i>		
<i>Centennial</i>	5	<i>purple</i>	3	<i>d</i>	<i>i</i>	
<i>Centennial Spirit</i>	12	<i>dark red</i>	2	<i>s</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Cherokee</i>	14	<i>red</i>	2	<i>s</i>	<i>i</i>	
<i>Chica Pink</i>	4	<i>bright pink</i>	1	<i>d</i>		
<i>Chica Red</i>	4	<i>rosy red</i>	1	<i>d</i>	<i>i</i>	
<i>Chickasaw</i>	3	<i>pink-lavender</i>	5	<i>m</i>	<i>h</i>	
<i>Chisam Fire</i>	5	<i>bright red</i>	1	<i>d-w</i>	<i>h</i>	
<i>Choctaw</i>	25	<i>bright pink</i>	4	<i>t</i>	<i>h</i>	<i>c-b</i>
<i>Christiana</i>	10	<i>dark red</i>	1	<i>s</i>	<i>i</i>	
<i>Christmas Time</i> (<i>Christmastime</i>)	20	<i>white</i>	3	<i>t</i>		
<i>Comanche</i>	18	<i>coral pink</i>	5	<i>t</i>	<i>h</i>	<i>c-b</i>
<i>Conestoga</i>	18	<i>pale lavender</i>	2	<i>t</i>	<i>i</i>	
<i>Cordon Bleu</i> (<i>Cordon Blue</i>) (<i>Improved Cordon Bleu</i>) (<i>Louisa</i>)	4	<i>lavender-blue</i>	1	<i>d-w</i>	<i>i</i>	
<i>Cotton Ball</i>	20	<i>white</i>	3	<i>t</i>	<i>i</i>	
<i>Country Red</i> (<i>Pure Red</i>) (<i>Durant Red</i>)	20	<i>dark red</i>	5	<i>t</i>	<i>i</i>	
<i>Creole</i>	3	<i>med. red</i>	1	<i>m-w</i>		
<i>Crimson</i> (<i>Carmine</i>) (<i>Carmine Pink</i>)	25	<i>deep red</i>	3	<i>t</i>		
<i>Dallas Red</i>	25	<i>red</i>	5	<i>t</i>	<i>i</i>	
<i>Deep Red</i>	20	<i>red</i>	0	<i>t</i>	<i>i</i>	
<i>Delta Blush</i> (<i>Pink Blush (old)</i>)	3	<i>pink</i>	1	<i>m-w</i>	<i>i</i>	
<i>Desal 173</i> (<i>Alienor</i>) (<i>Alienor d' Aquitaine</i>)	10	<i>dark pink</i>	1	<i>s-w</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Desand 081</i> (<i>Souvenir d'Andre Desmartis</i>)	25	red	3	<i>t</i>	<i>h</i>	
<i>Desber 102</i> (<i>Bergerac</i>) (<i>Dester 102</i>)	9	light red	3	<i>s</i>	<i>i</i>	
<i>Descha 14821</i> (<i>Peacharmant</i>)	18	light purple	3	<i>t</i>	<i>i</i>	
<i>Desemi 103</i> (<i>Saint Emilion</i>)	20	crimson red	3	<i>t</i>	<i>h</i>	
<i>Desjan 086</i> (<i>Jeanne Desmartis</i>)	18	rosy-pink	3	<i>t</i>	<i>i</i>	
<i>Desmon 104</i> (<i>Bengali</i>) (<i>Monbazillac</i>)	24	clear rose pink	3	<i>t</i>	<i>i</i>	
<i>Desmou 083</i> (<i>La Mousson</i>)	16	bright rose pink	3	<i>t</i>	<i>i</i>	
<i>Desper 14821</i> (<i>Perigord Pourpre</i>)	18	bright red	3	<i>t</i>	<i>i</i>	
<i>Desyan 184</i> (<i>Yang-Tse</i>)	25	crimson red	3	<i>t</i>	<i>i</i>	
<i>Diane</i>	20	lavender	3	<i>t</i>	<i>h</i>	
<i>Dwarf Blue</i> (<i>Baker Dwarf Blue</i>) (<i>Blue Midget</i>) (<i>Nana Blue</i>) (<i>Nana Corrulea</i>) (<i>Coerulea Nana</i>)	10	lavender-blue	2	<i>s</i>	<i>i</i>	
<i>Dwarf Crimson</i> (<i>Griffing's Dwarf Crimson</i>)	8	bright red	0	<i>d</i>	<i>i</i>	
<i>Dwarf Dark Red</i>	8	red	2	<i>d</i>	<i>i</i>	
<i>Dwarf Orchid</i>	5	med. orchid	2	<i>d</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Dwarf Purple</i> (<i>Dwarf Lavender</i>) (<i>Dwarf Lavender-Blue</i>) (<i>Frosty Blue</i>) (<i>Hardy Dwarf Lavender-Blue</i>) (<i>Lavendula Nana</i>) (<i>Lilac Nana</i>) (<i>Nana Lavendula</i>) (<i>Nana Purpurea</i>)	10	<i>rosy-purple</i>	3	<i>s</i>	<i>i</i>	
<i>Dwarf White</i>	6	<i>white</i>	1	<i>d</i>		
<i>Dynamite</i>	22	<i>cherry red</i>	3	<i>t</i>	<i>i</i>	
<i>Elfin</i>	5	<i>dark pink</i>	3	<i>d</i>		
<i>Far West</i>	20	<i>light pink</i>	2	<i>t</i>		
<i>Fantasy</i>	45	<i>white</i>	5	<i>t</i>	<i>f</i>	<i>c-b</i>
<i>Firebird</i> (<i>Firebird Red</i>)	26	<i>bright red</i>	2	<i>t</i>	<i>i</i>	
<i>Glendora White</i> (<i>Glendora</i>) (<i>Jetstream</i>)	25	<i>white</i>	3	<i>t</i>	<i>i</i>	
<i>Glowing Rose</i>	10	<i>pink</i>	3	<i>s</i>	<i>i</i>	
<i>Griffith Pink</i>	18	<i>pink</i>	4	<i>t</i>	<i>i</i>	
<i>Hardy Lavender</i>	30	<i>medium lavender</i>	3	<i>t</i>		
<i>Heavenly Blue</i>	5	<i>med. blue</i>	2	<i>d</i>	<i>h</i>	
<i>Hope</i>	5	<i>pinkish-white</i>	4	<i>d</i>	<i>i</i>	
<i>Hopi</i>	9	<i>medium pink</i>	5	<i>s</i>	<i>h</i>	<i>c-b</i>
<i>Houston</i>	3	<i>med. red</i>	1	<i>m-w</i>	<i>i</i>	
<i>Houston Red</i>	18	<i>dark magenta</i>	3	<i>t</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Ingleside Pink</i>	20	<i>pink</i>	3	<i>t</i>	<i>i</i>	
<i>Ingleside White</i>	26	<i>white</i>	3	<i>t</i>	<i>i</i>	
<i>Issai-Sarusubri</i>	10	<i>redish-purple</i>	3	<i>s</i>	<i>i</i>	
<i>Jet Stream</i> (<i>Glendora</i>)	25	<i>white</i>	3	<i>t</i>	<i>i</i>	
<i>Kiowa</i>	28	<i>white</i>	5	<i>t</i>	<i>f</i>	<i>c-b</i>
<i>Lancasteri</i>	14	<i>pale lavender</i>	3	<i>s</i>	<i>h</i>	
<i>Lafayette</i> (<i>Lavender Blush</i>)	3	<i>light lavender-white</i>	1	<i>m-w</i>	<i>i</i>	
<i>Lavender Beauty</i>	5	<i>lavender</i>	1	<i>d</i>	<i>i</i>	
<i>Lipan</i>	18	<i>purplish lavender</i>	5	<i>t</i>	<i>h</i>	<i>c-b</i>
<i>Little Red</i>	5	<i>red</i>	1	<i>d</i>		
<i>Little Silver</i>	5	<i>light lavender</i>	1	<i>d</i>	<i>i</i>	
<i>Low Flame</i> (<i>Dwarf Low Flame</i>) (<i>Flame</i>)	14	<i>red</i>	3	<i>s</i>	<i>i</i>	
<i>Magnifica</i>	5	<i>deep purple</i>	1	<i>d</i>		
<i>Maiden Blush</i>	3	<i>pink</i>	1	<i>m</i>	<i>h</i>	
<i>Mandi</i> (<i>Mandi Red</i>) (<i>Mandy</i>)	10	<i>dark red</i>	1	<i>s</i>		
<i>Mardi Gras</i>	3	<i>purple</i>	1	<i>m-w</i>	<i>i</i>	
<i>Mauvilla</i>	8	<i>red</i>	3	<i>d</i>	<i>i</i>	
<i>Miami</i>	25	<i>dark pink</i>	5	<i>t</i>	<i>h</i>	<i>b</i>
<i>Moned</i> (<i>Chica Red</i>)	4	<i>rosy-red</i>	3	<i>d</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Moners</i> (<i>Petite Embers</i>) (<i>Petite Emberg</i>)	5	<i>rosy-red</i>	2	<i>d</i>	<i>h</i>	<i>c</i>
<i>MONHID</i> (<i>Petite Orchid</i>)	5	<i>dark lavender</i>	1	<i>d</i>	<i>h</i>	<i>c</i>
<i>Monia</i> (<i>Majestic Orchid</i>) (<i>Majestic Cloud</i>) (<i>Tall Orchid</i>)	18	<i>lavender</i>	3	<i>t</i>	<i>h</i>	
<i>MONIMP</i> (<i>Petite Red Imp</i>) (<i>Petite Red Improved</i>)	5	<i>lavender</i>	0	<i>d</i>	<i>h</i>	
<i>Monink</i> (<i>Chica Pink</i>)	4	<i>bright pink</i>	2	<i>d</i>	<i>i</i>	
<i>Monkie</i> (<i>Petite Pinkie</i>)	5	<i>clear pink</i>	2	<i>d</i>	<i>h</i>	<i>c</i>
<i>Monow</i> (<i>Petite Snow</i>) (<i>Dwarf Snow</i>)	5	<i>bright white</i>	2	<i>d</i>	<i>h</i>	
<i>Monum</i> (<i>Petite Plum</i>)	5	<i>deep purple</i>	2	<i>d</i>	<i>h</i>	
<i>Munson</i>	24	<i>pink</i>	0	<i>t</i>	<i>i</i>	
<i>Muskogee</i> (<i>Muscogee</i>)	28	<i>light lavender</i>	5	<i>t</i>	<i>h</i>	<i>c</i>
<i>Natchez</i>	35	<i>white</i>	5	<i>t</i>	<i>h</i>	<i>c-b</i>
<i>Near East</i> (<i>Debaillon Pink</i>) (<i>Shell Pink</i>)	20	<i>light pink</i>	2	<i>t</i>	<i>i</i>	
<i>New Orleans</i> (<i>Passion</i>)	2	<i>purple</i>	1	<i>m-w</i>		

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
<i>New Pure White</i>	24	white	5 = best, 0 = poor 3	t	i	
<i>Nikki</i>	5	bi-color - pink/white	1	d		c
<i>Nivea</i>	9	white	3	s	i	
<i>Okmulgee</i> (<i>Ocmulgee</i>)	6	dark red	1	d		c
<i>Old Fashioned Arkansas</i>	18	redish-pink	3	t	i	
<i>Orbyn Atkins</i> (<i>Orbin Atkins</i>) (<i>Orbin Adkins</i>)	24	white	0	t		
<i>Orchid</i>	26	lavender	3	t	i	
<i>Osage</i>	18	clear pink	5	t-w	h	c-b
<i>Ozark Spring</i> (<i>Ozark Springs</i>) (<i>Ozard Spring</i>) (<i>Oklahoma Spring</i>)	5	light lavender	3	d	i	
<i>Parade Purple</i> (<i>Purple Parade</i>) (<i>Select Purple</i>)	25	purple	3	t	i	
<i>Pecos</i>	9	med. pink	5	s	h	
<i>Peppermint Lace</i>	12	pink with white	1	s	h	
<i>Petit' Canaille - Pink</i>	5	pink	1	d	i	
<i>Petit' Canaille - Red</i>	5	red	1	d	i	
<i>Petit' Canaille - Lavender</i>	5	lavender	1	d	i	
<i>Petit' Canaille - White</i>	5	white	1	d	i	
<i>Petit' Canaille - Rose</i>	5	pink	1	d	i	
<i>Petite Embers</i>	5	rose red	1	d		
<i>Petite Orchid</i>	5	dark orchid	1	d		
<i>Petite Pinkie</i>	5	clear pink	1	d		
<i>Petite Plum</i>	5	plum purple	1	d		
<i>Petite Red Imp</i>	5	crimson red	1	d		

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Petite Ruby</i> (<i>Petite Red Ruby</i>)	5	<i>deep ruby red</i>	1	<i>d</i>	<i>i</i>	
<i>Petite Snow</i>	5	<i>white</i>	1	<i>d</i>		
<i>Pink Blush (new)</i>	3	<i>light pink</i>	1	<i>m-w</i>	<i>i</i>	
<i>Pink Lace</i>	18	<i>bright pink</i>	3	<i>t</i>	<i>i</i>	
<i>Pink Parfait</i>	5	<i>clear pink</i>	1	<i>d</i>	<i>i</i>	
<i>Pink Ruffles</i> (<i>Pink Ruffle</i>) (<i>Dwarf Pink Ruffles</i>)	5	<i>pink</i>	2	<i>d</i>		
<i>Pink Snowflake</i>	5	<i>pink/white</i>	1	<i>d</i>		
<i>Pink Wonder</i>	5	<i>light pink</i>	1	<i>d</i>		
<i>Piroska</i>	24	<i>red</i>	2	<i>t</i>	<i>i</i>	
<i>Pixie White</i>	3	<i>white</i>	1	<i>m</i>		
<i>Pixie Pink</i>	4	<i>lavender-pink</i>	1	<i>d</i>		
<i>Pocomoke</i>	3	<i>deep rosy-pink</i>	4	<i>m-w</i>	<i>h</i>	
<i>Pomona Red</i>	18	<i>red</i>	3	<i>t</i>		
<i>Potomac</i>	24	<i>clear pink</i>	3	<i>t</i>	<i>i</i>	
<i>Powhatan</i>	18	<i>lavender</i>	3	<i>t</i>	<i>i</i>	
<i>Prairie Lace</i>	10	<i>pink / white</i>	2	<i>s</i>	<i>i</i>	
<i>Purity</i>	18	<i>white</i>	3	<i>t</i>		
<i>Purple Queen</i>	5	<i>reddish-purple</i>	1	<i>d</i>	<i>i</i>	
<i>Purple Rain</i>	5	<i>dark purple</i>	2	<i>d</i>	<i>i</i>	
<i>Purple Velvet</i>	5	<i>purple</i>	1	<i>d</i>	<i>i</i>	
<i>Purpurea</i> (<i>Purple</i>) (<i>Lavender</i>) (<i>Lavendula</i>) (<i>Lilac</i>) (<i>Lilac Purple</i>) (<i>Violacea</i>)	26	<i>bright purple</i>	3	<i>t</i>	<i>i</i>	
<i>Queens Lace</i>	18	<i>bi-color - light red/white</i>	3	<i>t</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Raspberry Sundae</i>	18	<i>pinkish-red / pale pink</i>	0	<i>t</i>	<i>i</i>	
<i>Red Imperator</i>	9	<i>red</i>	3	<i>s</i>	<i>i</i>	
<i>Red Flame</i>	5	<i>red</i>	1	<i>d</i>	<i>i</i>	
<i>Red Rocket</i>	25	<i>cherry red</i>	4	<i>t</i>		
<i>Red Snowflake</i>	18	<i>dark red / white</i>	3	<i>t</i>	<i>i</i>	
<i>Red Sport</i>	18	<i>red</i>	3	<i>t</i>	<i>i</i>	
<i>Regal Red</i> (<i>Byers Regal Red</i>)	18	<i>dark red</i>	4	<i>t</i>	<i>i</i>	
<i>Regina</i> (<i>Regia</i>)	24	<i>red</i>	3	<i>t</i>	<i>i</i>	
<i>Reginae Orchid</i>	18	<i>light lavender</i>	2	<i>t</i>	<i>h</i>	
<i>Rhapsody</i>	5	<i>red</i>	3	<i>d</i>	<i>i</i>	
<i>Rosa</i>	26	<i>rosy-pink</i>	3	<i>t</i>	<i>h</i>	
<i>Rosea</i> (<i>Rose</i>) (<i>Rosea Elegans</i>)	24	<i>deep pink</i>	1	<i>t</i>	<i>i</i>	
<i>Rosea Nova</i>	24	<i>pink</i>	3	<i>t</i>		
<i>Royal Pink</i>	18	<i>pink</i>	3	<i>t</i>	<i>h</i>	<i>c</i>
<i>Royal Purple</i>	18	<i>purple</i>	3	<i>t</i>	<i>i</i>	
<i>Royal Velvet</i>	18	<i>bright pink</i>	3	<i>t</i>	<i>i</i>	
<i>Royalty</i> (<i>Dwarf Royalty</i>)	5	<i>purple</i>	1	<i>d</i>	<i>i</i>	
<i>Rubra</i> (<i>Red</i>) (<i>Magenta</i>) (<i>Rubis</i>) (<i>Rubrum</i>)	9	<i>crimson red</i>	3	<i>s</i>	<i>i</i>	
<i>Rubra Compacta</i> (<i>Nana Rubra</i>) (<i>Dwarf Red</i>)	9	<i>red</i>	3	<i>d</i>	<i>i</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
<i>Rubra Grayi</i> (Gray's Red) (Grey's Red)	24	red	<small>5 = best, 0 = poor</small> 0	<i>t</i>		
<i>Sacramento</i>	3	deep red	1	<i>m</i>	<i>i</i>	
<i>Saharanpur</i>	24	dark pink	3	<i>t</i>	<i>h</i>	
<i>Sarah's Favorite</i> (5570) (Sarah's Hardy White) (Dood's White) (Hybrid White)	26	white	5	<i>t</i>	<i>h</i>	<i>c</i>
<i>Scarlet</i>	16	red	0	<i>t</i>		
<i>Scarlet Gem</i>	18	red	3	<i>t</i>	<i>i</i>	
<i>Seminole</i>	18	clear redish-pink	2	<i>t</i>	<i>i</i>	
<i>Sioux</i>	18	dark pink	5	<i>t</i>	<i>h</i>	<i>c-b</i>
<i>Snow</i>	6	white	2	<i>d</i>	<i>i</i>	
<i>Snowbaby</i> (New Snow Baby) (Dwarf Snow Baby) (Dwarf White) (Frosty Snow) (Alba Nana)	5	white	1	<i>d</i>	<i>i</i>	
<i>Snow Lace</i>	18	white	3	<i>t-w</i>	<i>i</i>	
<i>Southern Belle</i>	18	flesh pink	2	<i>t</i>	<i>i</i>	
<i>Special Red</i>	25	coral red	1	<i>t</i>	<i>i</i>	
<i>Spice Pink</i>	18	pinkish-white	0	<i>t</i>	<i>i</i>	
<i>St. Michael's White</i>	30	white	3	<i>t</i>	<i>i</i>	
<i>Summer and Summer</i>	2	purple	2	<i>m</i>	<i>i</i>	
<i>Summer Wine</i>	5	deep pink	2	<i>d</i>	<i>i</i>	
<i>Tawakoni Pink</i>	18	med. hot pink	3	<i>t</i>		
<i>The Bride</i>	24	light pink	0	<i>t</i>	<i>h</i>	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
<i>Tightwad Red</i>	4	light red	5 = best, 0 = poor 5	d		
<i>Tinsley</i>	25	lavender	3	t	h	
<i>Tiny Fire</i>	5	red	1	d	i	
<i>Tom Dodd</i>	5	orange red	3	d	i	
<i>Tonto</i>	10	bright pinkish-red	5	s	h	c
<i>Townhouse</i>	25	white	5	t	f	c-b
<i>True Red</i>	9	rose red	3	s	i	
<i>Tuscaloosa Red</i>	26	red	2	t	i	
<i>Tuscarora</i>	24	dark coral pink	5	t	h	
<i>Tuskegee</i>	20	dark pink	5	t	h	c-b
<i>Twilight</i>	20	dark purple	1	t	i	
<i>Velma's Royal Delight</i>	5	magenta	4	d	i	
<i>Victor</i> (Victor Dwarf)	5	deep red	3	d	i	
<i>Watermelon Red</i>	25	bright red	1	t	i	
<i>Weeping Lavender</i>	5	lavender	1	d-w	i	
<i>Weeping Pink</i> (Pendula) (Prostrata Griffin) (Prostrata) (Rosea Pendula)	18	rosy-pink	3	t-w	i	
<i>Weeping Red</i>	5	red	1	d-w	i	
<i>Weeping White</i>	18	white	3	t-w	i	
<i>Whit I</i> (Raspberry Sundae)	18	bi-color-pinkish-red/pale-pink	0	t	i	
<i>Whit II</i> (Dynamite)	22	cherry red	3	t	i	
<i>Whit III</i> (Royal Velvet) (Pink Velour)	18	bright pink	3	t	i	

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = L. indica; f = L. fauriei; h = hybrid; B = great bark; C = best relative cold hardiness

crapemyrtle cultivar (other names)	height (ft.)	flower color	disease resistance	form	genetics	features
			<i>5 = best, 0 = poor</i>			
<i>Whit IV</i> (<i>Red Rocket</i>)	26	<i>bright red</i>	3	<i>t</i>	<i>i</i>	
<i>Whit V</i> (<i>Tightwad</i>) (<i>Firecracker</i>)	5	<i>light red</i>	4	<i>d</i>	<i>i</i>	
<i>White Cloud</i> (<i>Cloud White</i>)	24	<i>clear white</i>	3	<i>t</i>		
<i>White Layers</i>	9	<i>white</i>	2	<i>s~w</i>	<i>i</i>	
<i>Wichita</i>	22	<i>pinkish-lavender</i>	5	<i>t</i>	<i>h</i>	<i>c-b</i>
<i>Wm. Toovey</i> (<i>William Toovey</i>) (<i>Rubra Nana</i>)	18	<i>pinkish red</i>	3	<i>t</i>	<i>i</i>	
<i>Worlds Fair</i>	3	<i>deep red</i>	1	<i>m-w</i>	<i>i</i>	
<i>Yuma</i>	18	<i>double med. lavender</i>	4	<i>t</i>	<i>h</i>	
<i>Yvonne</i>	15	<i>pink-lavender</i>	3	<i>t</i>	<i>h</i>	
<i>Zuni</i>	10	<i>med. lavender</i>	5	<i>s</i>	<i>h</i>	
<div style="border: 1px solid black; padding: 10px;"> <p>Primary Sources of Information: Experiment Station and Extension Service WEB sites of Auburn, Clemson, Florida, and the University of Georgia. The Crape Myrtle (<i>Lagerstroemia</i> spp.) site of Texas A & M = [dallas.tamu.edu/woody/cmirtle] Characteristics of Crape Myrtle Varieties site of Texas A & M, PLANTanswers, Aggie Horticulture = [aggie-horticulture.tamu.edu]</p> </div>						

w = weeping; m = miniature (<3 feet tall); d = dwarf (4-8 feet); s = standard (9-15 feet); t = tall (>16 feet); i = *L. indica*; f = *L. fauriei*; h = hybrid; B = great bark; C = best relative cold hardiness