

Pest Review Bibliography For Loblolly Pine (*Pinus taeda*)

by Dr. Kim D. Coder, University of Georgia

Sept. 1997

The following research literature citations are from approximately the last 15 years. These are selected citations concerning arthropods, diseases, nematodes, and general pests of loblolly pine (*Pinus taeda*). This publication lists citations considered to be significant additions to the pest and host information base. This publication is intended to assist plant health care (PHC) practitioners.

Adams DE, Lenhart JD, Vaughn AB, Lapongan J. 1996. Predicting survival of east Texas loblolly and slash pine plantations infected with fusiform rust. *Southern Journal of Applied Forestry*. 20:30-35

Anderson RF, Doggett CA. 1980. Some relationships between infestations by the southern pine beetle (*Dendroctonus frontalis*) and stand conditions. Forestry Note Division of Forest Resources, North Carolina. No.49, 16 pp. Raleigh, North Carolina, USA

Anderson RF, Doggett CA. 1993. Host preference of southern pine beetle in North Carolina. Forestry Note Division of Forest Resources, North Carolina. 1993, No.66, 7 pp. Raleigh, North Carolina, USA

Anderson RL, McCartney TC, Cost ND, Devine H, Botkin M. 1988. Fusiform rust hazard maps for loblolly and slash pines. Research Note Southeastern Forest Experiment Station, USDA Forest Service. SE-351, 7 pp.

Anderson RL, McClure JP, Cost N, Uhler RJ. 1986. Estimating fusiform rust losses in five southeast states. *Southern Journal of Applied Forestry*. 10:237-240.

Anderson RL, Walkinshaw CH. 1989. Correlation of greenhouse inoculations of longleaf pine with *Cronartium quercuum* f.sp. fusiform with field ranking of resistance. *Plant Disease*. 73:859

Anderson WC, Guldin RW, Vasievich JM. 1987. Assessing the risk of insect attack in plantation investments. *Journal of Forestry*. 85:46-47.

Anderson WC, Guldin RW, Vasievich JM. 1987. Assessing risk to plantation investments from insect attacks. Research Paper Southern Forest Experiment Station, USDA Forest Service. SO-231, 6 pp.

Baker FA, Verbyla DL, Hodges CS Jr., Ross EW. 1993. Classification and regression tree analysis for assessing hazard of pine mortality caused by *Heterobasidion annosum*. *Plant Disease*. 77:136-139

Barnard EI, Gilly SP, Ash EC. 1994. An evaluation of dazomet and metam sodium soil fumigants for control of *Macrophomina phaseolina* in a Florida forest nursery. *Tree Planters' Notes*. 45:91-95

Barnard EL. 1994. Nursery to field carryover and post outplanting impact of *Macrophomina phaseolina* on loblolly pine on a cutover forest site in North Central Florida. *Tree Planters' Notes*. 45:68-71



The University of Georgia

THE UNIVERSITY OF GEORGIA, THE UNITED STATES DEPARTMENT OF AGRICULTURE
AND COUNTIES OF THE STATE COOPERATING THE COOPERATIVE EXTENSION
SERVICE OFFERS EDUCATIONAL PROGRAMS, ASSISTANCE AND MATERIALS TO ALL
PEOPLE WITHOUT REGARD TO RACE, COLOR, NATIONAL ORIGIN,
AGE, SEX OR HANDICAP STATUS.
A UNIT OF THE UNIVERSITY SYSTEM OF GEORGIA.
AN EQUAL OPPORTUNITY/AFFIRMATIVE ACTION ORGANIZATION

University of Georgia Cooperative Extension Service publication FOR97-25

- Barrows Broaddus J. 1990. Colonization of clones and seed of loblolly pine following inoculation with Fusarium subglutinans. Plant Disease. 74:1002-1005
- Berisford CW, Brady UK, Fatzinger CW, Ebel BH. 1986. Evaluation of a repellent for prevention of attacks by three species of southern pine bark beetles. Journal of Entomological Science. 21: 316-318.
- Berisford CW, Godbee JF Jr., Ross DW. 1989. Impact of pine tip moth control, weed control, and fertilizer on growth and form of loblolly pine. Insects affecting reforestation: biology and damage. Pp.130-136. Forestry Canada, Pacific and Yukon Region, Victoria, Canada
- Bramlett DL. 1987. Protection of pine seed orchards in the southeastern United States. Forest Ecology and Management. 19:199-208. Flowering and seed bearing in forest seed orchards. Proceedings of the International Symposium. IUFRO S2.01.00 and 2.01.05. Sept. 1985, Kornik, Poland.
- Bridges JR. 1987. Effects of terpenoid compounds on growth of symbiotic fungi associated with the southern pine beetle. Phytopathology.77: 83-85.
- Bridges JR, Perry TJ. 1987. Ceratocystiopsis ranaculosus sp. nov. associated with the southern pine beetle. Mycologia. 79:630-633.
- Brissette JC, Barnett JP, Jones JP. 1996. Fungicides improve field performance of stored loblolly and longleaf pine seedlings. Southern Journal of Applied Forestry. 20:5-9
- Brown MW, Nebeker TE, Honea CR. 1987. Thinning increases loblolly pine vigor and resistance to bark beetles. Southern Journal of Applied Forestry. 11:28-31
- Burkhart HE, Munsey DE Jr., Amateis RL, Hedden RL. 1992. Predicting impact of southern pine beetle infestations on rotation age yield of loblolly pine stands. Forest Ecology and Management. 47:261-268.
- Byrne SV, Wentworth TR, Nusser SM. 1987. A moisture strain index for loblolly pine. Canadian Journal of Forest Research. 17:23-26
- Cade SC, Hedden RL. 1987. Growth impact of pine tip moth on loblolly pine plantations in the Ouachita Mountains of Arkansas. Southern Journal of Applied Forestry. 11:128-133
- Cade SC, Hedden RL. 1989. Evidence for tip moth (Rhyacionia frustrana) resistance in loblolly pine (Pinus taeda). Insects affecting reforestation: biology and damage. pp.137-144. Forestry Canada, Pacific and Yukon Region, Victoria, Canada
- Cameron RS. 1989. Promising new pesticides for cone and seed insect control in the southern United States. Proceedings of the 3rd Cone and Seed Insects Working Party Conference, held in Victoria, British Columbia, Canada, on 26-30 June 1988. Pp.193-202. Pacific Forestry Centre, Forestry Canada, Victoria, British Columbia, Canada
- Cameron RS, Billings RF. 1988. Southern pine beetle: Factors associated with spot occurrence and spread in young plantations. Southern Journal of Applied Forestry. 12:208-214
- Cameron RS, DeBarr GL. 1989. Responses of Dioryctria clarioralis, D.disclusa and D.merkeli to pheromone baits in Georgia and Texas. Proceedings of the 3rd Cone and Seed Insects Working Party Conference, held in Victoria, British Columbia, Canada, on 26-30 June 1988. Pp.174-180. Pacific Forestry Centre, Forestry Canada, Victoria, British Columbia, Canada
- Carey WA, Keller WD. 1996. Eight year field performance of pine seedlings after mycorrhizae were triadimefon inhibited in the nursery. Southern Journal of Applied Forestry. 20:42-44
- Carey WA, Kelley WD. 1993. Seedling production trends and fusiform rust control practices at southern nurseries, 1981-1991. Southern Journal of Applied Forestry. 17:207-211
- Carey WA, Kelley WD. 1994. Interaction of ozone exposure and Fusarium subglutinans inoculation on growth and disease development of loblolly pine seedlings. Environmental Pollution. 84:35-43

- Carson SD, Young CH. 1987. Effect of inoculum density and fertilization on greenhouse screening of loblolly pine seedlings for resistance to fusiform rust. *Phytopathology*. 77:1186-1191.
- Cerrano Moreira AF, Berisford CW, Hedden RL, Silva JAA da, Da Silva JAA. 1994. Methodology for rating *Pinus taeda* stands for pine tip moth, *Rhyacionia* spp. infestation. *Anais da Sociedade Entomologica do Brasil*. 23:227-236
- Chung DY, Kwon HM, Tak WS, Kim JH, Kim TS. 1984. Control of cone insects of four pine species by soil application of furadan (carbofuran) at two locations. Research Report of the Institute of Forest Genetics, Korea Republic. No.20, 70-73.
- Clarke SR, DeBarr GL, Berisford CW. 1988. Differential susceptibility of *Toumeyella pini* to pyrethroid and organophosphate insecticides: a factor in outbreaks in southern pine seed orchards. *Journal of Economic Entomology*. 81:1443-1445.
- Clarke SR, DeBarr GL, Berisford CW. 1989. The life history of *Toumeyella pini* in loblolly pine seed orchards in Georgia. *Canadian Entomologist*. 121:853-860.
- Clarke SR, DeBarr GL, Berisford CW. 1989. Life history of the woolly pine scale *Pseudophilippia quaintancii* in loblolly pine seed orchards. *Journal of Entomological Science*. 24:365-372
- Clarke SR, DeBarr GL, Berisford CW. 1990. Effects of fenvalerate and azinphosmethyl on scale insects and their natural enemies in loblolly pine seed orchards. Research Paper Southeastern Forest Experiment Station, USDA Forest Service. SE-279.
- Clarke SR, DeBarr GL, Berisford CW. 1990. Life history of *Oracella acute* in loblolly pine seed orchards in Georgia. *Environmental Entomology*. 19:99-103.
- Clarke SR, Negron JF, DeBarr GL. 1992. Effects of four pyrethroids on scale insect (*Homoptera*) populations and their natural enemies in loblolly and shortleaf pine seed orchards. *Journal of Economic Entomology*. 85:1246-1252.
- Clarke SR, Stephen FM, Dunn JE. 1990. Spatial distribution of the Nantucket pine tip moth in newly established loblolly pine plantations. *Journal of Entomological Science*. 25: 209-215.
- Conner RN, Rudolph DC, Kulhavy DL, Snow AE. 1991. Causes of mortality of red cockaded woodpecker cavity trees. *Journal of Wildlife Management*. 55:531-537
- Cook SP, Hain FP. 1987. Four parameters of the wound response of loblolly and shortleaf pines to inoculation with the blue staining fungus associated with the southern pine beetle. *Canadian Journal of Botany*. 65:2403-2409
- Cook SP, Hain FP. 1987. Four parameters of the wound response of loblolly and shortleaf pines to inoculation with the blue staining fungus associated with the southern pine beetle. *Canadian Journal of Botany*. 65:2403-2409.
- Cook SP, Hain FP. 1987. Susceptibility of trees to southern pine beetle, *Dendroctonus frontalis*. *Environmental Entomology*. 16:9-14.
- Cook SP, Hain FP. 1988. Toxicity of host monoterpenes to *Dendroctonus frontalis* and *Ips calligraphus*. *Journal of Entomological Science*. 23:287-292.
- Cook SP, Hain FP. 1988. Wound response of loblolly and shortleaf pine attacked or reattacked by *Dendroctonus frontalis* or its fungal associate, *Ceratocystis minor*. *Canadian Journal of Forest Research*. 18:33-37
- Cook SP, Hain FP, Nappen PB. 1986. Seasonality of the hypersensitive response by loblolly and shortleaf pine to inoculation with a fungal associate of the southern pine beetle. *Journal of Entomological Science*. 21:283-285
- Cook SP, Hain FP, Smith HR. 1994. Oviposition and pupal survival of gypsy moth in Virginia and North Carolina pine hardwood forests. *Environmental Entomology*. 23:360-366
- Cormier KL, Reich RM, Czaplowski RL, Bechtold WA. 1992. Evaluation of weighted regression and sample size in developing a taper model for loblolly pine. *Forest Ecology and Management*. 53:65-76

- DeAngelis JD, Nebeker TE, Hodges JD. 1986. Influence of tree age and growth rate on the radial resin duct system in loblolly pine (Pinus taeda). Canadian Journal of Botany. 64:1046-1049.
- Dixon WN. 1987. Nantucket pine tip moth, Rhyacionia frustrana. Entomology Circular, Division of Plant Industry, Florida Department of Agriculture and Consumer Services. No.298, 2pp.
- Doggett C, Holzman S, Rogers D, Smith J, Barron H. 1991. Survey of fusiform rust on loblolly pine in North Carolina. Forestry Note Division of Forest Resources, North Carolina. No.64, 10 pp. Raleigh, North Carolina, USA
- Doggett C, Rogers D, Smith J, Barron H. 1991. Within tree distribution of fusiform rust on loblolly pine. Forestry Note Division of Forest Resources, North Carolina. No.63, 14 pp. Raleigh, North Carolina, USA
- Doggett CA, Grady CR, Green HJ, Kunselman MB, Layman H, Taylor S. 1977. Seedling debarking weevils in North Carolina. Forestry Note Division of Forest Resources, North Carolina. No.31, 15 pp. Raleigh, North Carolina, USA
- Dunn JP, Lorio PL Jr.. 1992. Effects of bark girdling on carbohydrate supply and resistance of loblolly pine to southern pine beetle (Dendroctonus frontalis) attack. Forest Ecology and Management. 50:317-330
- Dunn JP, Lorio PL Jr.. 1993. Modified water regimes affect photosynthesis, xylem water potential, cambial growth, and resistance of juvenile Pinus taeda to Dendroctonus frontalis. Environmental Entomology. 22:948-957.
- Dwinell LD. 1986. Ecology of the pinewood nematode in southern pine chip piles. Research Paper, Southeastern Forest Experiment Station, USDA Forest Service. SE-258, 14 pp.
- Edwards MB, Dougherty PM. 1988. Controlling bermudagrass to establish loblolly pines. Proceedings of the 41st Annual Meeting of the Southern Weed Science Society. Pp.202-209, 3 ref.
- Esher RJ, Marx DH, Ursic SJ, Baker RL, Brown LR, Coleman DC. 1992. Simulated acid rain effects on fine roots, ectomycorrhizae, microorganisms, and invertebrates in pine forests of the southern United States. Water, Air, and Soil Pollution. 161:269-278.
- Fatzinger CW, Muse HD, Miller T, Bhattacharyya H. 1990. Survey and pest monitoring system for southern pine seed orchards. Southern Journal of Applied Forestry. 14:147-159.
- Fatzinger CW, Yates HO III, Hammond WJ, Hutto R. 1985. Insect caused cone and seed losses in treated and untreated loblolly pine seed orchards. Research Note, Southeastern Forest Experiment Station, USDA Forest Service. SE-333.
- Feaver MN. 1985. Grasshopper damage to pine seedlings at night in a seed orchard. Florida Entomologist. 68:694-696.
- Flamm RO, Cook SP, Wagner TL, Pulley PE, Coulson RN. 1987. Reemergence and emergence of Ips avulsus and Ips calligraphus. Environmental Entomology. 16:869-876.
- Flamm RO, Coulson RN, Beckley P, Pulley PE, Wagner TL. 1989. Maintenance of a phloem inhabiting guild. Environmental Entomology. 18:381-387
- Flamm RO, Pulley PE, Coulson RN. 1993. Colonization of disturbed trees by the southern pine bark beetle guild. Environmental Entomology. 22:62-70.
- Flamm RO, Wagner TL, Cook SP, Pulley PE, Coulson RN, McArdle TM. 1987. Host colonization by cohabiting Dendroctonus frontalis, Ips avulsus and I. calligraphus. Environmental Entomology. 16:390-399
- Foltz JL, Blakeslee GM. 1989. Insects associated with the intensive culture of Pinus elliottii and P.taeda in Florida. Insects affecting reforestation: biology and damage. Pp.19-26. Forestry Canada, Pacific and Yukon Region, Victoria, Canada
- Fraedrich SW. 1988. The influence of rhizosphere oxygen concentration and host nutrition on the susceptibility of shortleaf and loblolly pine roots to infection by Phytophthora cinnamomi. Dissertation Abstracts International. B, Sciences and Engineering. 49:16 B, Thesis, Clemson University, USA (1987), 169 pp.

- Fredericksen TS, Hedden RL, Williams SA. 1995. Susceptibility of loblolly pine to bark beetle attack following simulated wind stress. *Forest Ecology and Management*. 76:95-107.
- Froelich RC. 1989. Annual variation in fusiform rust infection of slash and loblolly pine seed lots in time replicated plantings. *Canadian Journal of Forest Research*. 19:1531-1537.
- Ge JJ, Cheng HR. 1993. Study on the resistance of three conifers to pine wood nematode. *Plant Quarantine Shanghai*. 7:337.
- Geron CD, Hafley WL. 1988. Impact of fusiform rust on product yields of loblolly pine plantations. *Southern Journal of Applied Forestry*. 12:226-231.
- Goldammer DS, Stephen FM, Paine TD. 1989. Average radial growth rate and chlamydospore production of *Ceratocystis minor*, *Ceratocystis minor* var. *barrasii*, and SJB 122 in culture. *Canadian Journal of Botany*. 67:3498-3505.
- Gopo JM, Katerere Y, Sibindi F. 1989. Isolation, purification and identification of mycoplasma like organisms as causal agent for the yellows disease in *Pinus* and in *Callitris* cypress in Zimbabwe. *Discovery and Innovation*. 1:82-86.
- Guldin RW. 1985. Pine release in unevenly stocked stands on droughty soils may be uneconomical. Research Paper Southern Forest Experiment Station, USDA Forest Service. SO-216.
- Hamelin RC, Doudrick RL, Nance WL. 1994. Genetic diversity in *Cronartium quercuum* f.sp *fusiforme* on loblolly pines in southern U.S. *Current Genetics*. 26:359-363.
- Harms WR, DeBell DS, Whitesell CD. 1994. Stand and tree characteristics and stockability in *Pinus taeda* plantations in Hawaii and South Carolina. *Canadian Journal of Forest Research*. 24:511-521
- Hayes JL, Ingram LL Jr., Strom BL, Roton LM, Boyette MW, Walsh MT, Vozzo JA. 1994. Identification of a host compound and its practical applications: 4 allylanisole as a bark beetle repellent. Research and applications of chemical sciences in forestry. Proceedings of the 4th Southern Station Chemical Sciences Meeting, February 1-2, 1994, Starkville, MS. General Technical Report Southern Forest Experiment Station, USDA Forest Service. SO-104, pp.69-79. New Orleans, Louisiana, USA
- Hayes JL, Meeker JR, Foltz JL, Strom BL. 1996. Suppression of bark beetles and protection of pines in the urban environment: a case study. *Journal of Arboriculture*. 22:67-74
- Hayes JL, Strom BL, Roton LM, Ingram LL Jr.. 1994. Repellent properties of the host compound 4 allylanisole to the southern pine beetle. *Journal of Chemical Ecology*. 20:1595-1615.
- Hedden RL, Belanger RP, Powers HR, Miller T. 1991. Relation of Nantucket pine tip moth attack and fusiform rust infection in loblolly pine families. *Southern Journal of Applied Forestry*. 15:204-208.
- Heikkinen HJ, Scheckler SE, Egan PJJ Jr., Williams CB Jr.. 1986. Incomplete abscission of needle clusters and resin release from artificially water stressed loblolly pine (*Pinus taeda*): a component for plant animal interactions. *American Journal of Botany*. 73:1384-1392.
- Hodge GR, White TL, Schmidt RA, Allen JE. 1993. Stability of rust infection ratios for resistant and susceptible slash and loblolly pine across rust hazard levels. *Southern Journal of Applied Forestry*. 17:188-192.
- Hood WM, Hedden RL, Berisford CW. 1988. Hazard rating forest sites for pine tip moth, *Rhyacionia* spp. in the upper Piedmont plateau. *Forest Science*. 34:1083-1093.
- Isagi Y. 1987. Relationships between the light conditions of stands and the defoliation of deciduous broadleaved trees by two leaf rolling weevil species, *Deporaus unicolor* and *Apoderus erythrogaster*. *Journal of the Japanese Forestry Society*. 69:482-486.
- Iwahori H, Futai K. 1996. Changes in nematode population and movement in pine seedlings with the development of pine wilt disease suppressive effect of water extracts from infected pine seedlings on the movement of nematodes. *Applied Entomology and Zoology*. 31:11-20

- Jacobi JC, Tainter FH, Oak SW. 1988. The effect of drought on growth decline of loblolly pine on littleleaf sites. *Plant Disease*. 72:294-297.
- Jang JC, Tainter FH. 1990. Cellular responses of pine callus to infection by *Phytophthora cinnamomi*. *Phytopathology*. 80:1347-1352.
- Jang JC, Tainter FH. 1991. Optimum tissue culture conditions for selection of resistance to *Phytophthora cinnamomi* in pine callus tissue. *Plant Cell Reports*. 9:488-491
- Jett JB, Guinness WM. 1992. Growth and wood properties in a Carolina sandhills pine seed source study. *Southern Journal of Applied Forestry*. 16:164-169
- Jewell FF Sr.. 1993. Histopathology of loblolly pine needles infected by *Mycosphaerella dearnessii*, the brown spot needle disease pathogen. *European Journal of Forest Pathology*. 23:110-119
- Kelley WD. 1988. Effect of short term storage of triadimefon treated loblolly pine seed on incidence of fusiform rust. *Southern Journal of Applied Forestry*. 12:18-20
- Kelly JM, Taylor GE Jr., Edwards NT, Adams MB, Edwards GS, Friend AL. 1993. Growth, physiology, and nutrition of loblolly pine seedlings stressed by ozone and acidic precipitation: a summary of the ROPIS South project. *Water, Air, and Soil Pollution*. 69:363-391
- Kinn DN. 1986. Incidence of the pinewood nematode in a southern pine beetle infestation in central Louisiana. *Journal of Entomological Science*. 21:114-117
- Kinn DN. 1987. Incidence of pinewood nematode dauerlarvae and phoretic mites associated with long horned beetles in central Louisiana. *Canadian Journal of Forest Research*. 17:187-190
- Ko JH, La YJ. 1985. Proceedings of the joint conference of IUFRO working parties on forest gall midges and rusts of pines. 1985. 171pp. Korean Forestry Society, Suweon, Korea.
- Kraus JF. 1986. Breeding shortleaf X loblolly pine hybrids for the development of fusiform rust resistant loblolly pine. *Southern Journal of Applied Forestry*. 10:195-197.
- Kuhlman EG. 1987. Effects of inoculation treatment with *Fusarium moniliforme* var. *subglutinans* on dieback of loblolly and slash pine seedlings. *Plant Disease*. 71:161-162.
- Kuhlman EG. 1988. Histology and progression of fusiform rust symptoms on inoculated loblolly pine seedlings. *Plant Disease*. 72:719-721
- Kuhlman EG. 1990. Frequency of single gall isolates of *Cronartium quercuum* f.sp. *fusiforme* with virulence toward three resistant loblolly pine families. *Phytopathology*. 80:614-617
- Kuhlman EG. 1992. Interaction of virulent single gall rust isolates of *Cronartium quercuum* f.sp. *fusiforme* and resistant families of loblolly pine. *Forest Science*. 38:641- 651.
- Kuhlman KG, Powers HR Jr.. 1988. Resistance response in half sib loblolly pine progenies after inoculation with *Cronartium quercuum* f.sp. *fusiforme*. *Phytopathology*. 78:484-487
- Kuhlman KG, Powers HR, Pepper WD. 1990. Resistant responses in loblolly pine to spray and injection inoculations with *Cronartium quercuum* f.sp. *fusiforme*. *Plant Disease*. 74:1013-1015
- Kuhlman KG, Powers HR, Pepper WD. 1995. Relative fusiform rust resistance of loblolly and slash pine sources and families in Georgia and South Carolina. Research Paper Southern Research Station, USDA Forest Service. SE-291
- Kuroda K, Yamada T, Ito S. 1991. *Bursaphelenchus xylophilus* induced pine wilt: factors associated with resistance. *European Journal of Forest Pathology*. 21:430-438

- Lantagne DO, Burger JA. 1988. Effect of site preparation intensity on the early growth of loblolly pine (Pinus taeda) and the incidence of pine tip moth (Rhyacionia spp.). *New Forests*. 2:219-229
- Lawrence LK. 1975. Relationship between number of pales weevils trapped and subsequent seedling mortality. *Forestry Note Division of Forest Resources, North Carolina*. No.23. Raleigh, North Carolina, USA
- Lee KiYoung, Lee ChanYoung, Lee KY, Lee CY. 1994. A study on the cone insects of some Pinus. *Research Bulletin of the Experiment Forests, Kangweon National University*. 14:139-151
- Lee SG, Lee KJ, La YJ, Yang SI, Yi CK. 1989. Identification of Lophodermium species associated with needle cast disease of pines in Korea. *Journal of Korean Forestry Society*. 78:218-227.
- Lenhart JD, Gregoire TG, Kronrad GD, Holley AG. 1994. Characterizing fusiform rust incidence and distribution in east Texas. *Southern Journal of Applied Forestry*. 18:29-34.
- Lenhart JD, Kronard GD, Fountain MS. 1993. Comparison of planted loblolly and slash pine performance in southeast Texas. *Southern Journal of Applied Forestry*. 17:26-31.
- Liebholt AM, Elmes GA, Halverson JA, Quimby J. 1994. Landscape characterization of forest susceptibility to gypsy moth defoliation. *Forest Science*. 40:18-29
- Lin ZS, Liang ZC. 1988. Comparative resistance of several pine species to brown spot needle blight and the origin of the disease. *Scientia Silvae Sinicae*. 24:25-47.
- Linhart YB. 1991. Disease, parasitism and herbivory: multidimensional challenges in plant evolution. *Trends in Ecology and Evolution*. 6:392-396
- Linit MJ, Kinn DN. 1993. Influence of pinewood nematode infection on the preformed defensive response of loblolly pine. *Environmental Entomology*. 22:1285-1293.
- Lorio PL Jr., Sommers RA. 1986. Evidence of competition for photosynthates between growth processes and oleoresin synthesis in Pinus taeda L. *Tree Physiology*. 2:301-306.
- Lorio PL Jr., Stephen FM, Paine TD. 1995. Environment and ontogeny modify loblolly pine response to induced acute water deficits and bark beetle attack. *Forest Ecology and Management*. 73:97-110
- Lovelady CN, Pulley PE, Coulson RN, Flamm RO. 1991. Relation of lightning to herbivory by the southern pine bark beetle guild. *Environmental Entomology*. 20:1279-1284.
- Lundquist JE. 1993. Spatial and temporal characteristics of canopy gaps caused by Armillaria root disease and their management implications in lowveld forests of South Africa. *European Journal of Forest Pathology*. 23:362-371.
- MacFall JS, Spaine P, Doudrick R, Johnson GA. 1994. Alterations in growth and water transport processes in fusiform rust galls of pine, determined by magnetic resonance microscopy. *Phytopathology*. 84:288-293
- Marx DH, Cordell CE. 1987. Triadimefon affects Pisolithus ectomycorrhizal development, fusiform rust, and growth of loblolly and slash pines in nurseries. *Research Paper, Southeastern Forest Experiment Station, USDA Forest Service*. SE-267. 14 pp.
- Matson PA, Hain FP, Mawby W. 1987. Indices of tree susceptibility to bark beetles vary with silvicultural treatment in a loblolly pine plantation. *Forest Ecology and Management*. 22:107-118
- McNab WH, Miller T, Brender EV. 1990. Growth and fusiform rust responses of Piedmont loblolly pine after several site preparation and regeneration methods. *Southern Journal of Applied Forestry*. 14:18-24
- Meeker JR, Kulhavy DL. 1992. Pine tip moth infestation rates as influenced by site and stand characteristics in loblolly pine plantations in east Texas. *Environmental Entomology*. 21:534-541

- Michelozzi M, White TL, Squillace AK, Lowe WJ. 1995. Monoterpene composition and fusiform rust resistance in slash and loblolly pines. *Canadian Journal of Forest Research*. 25:193-197
- Miller MC. 1986. Within tree effects of bark beetle insect associates on the emergence of Ips calligraphus. *Environmental Entomology*. 15:1104-1108.
- Miller MC, Kinn DN, Parresol BR. 1995. Effect of sodium N methylthiocarbamate with dimethyl sulfoxide on southern pine beetle development: results of initial field tests. *Journal of Applied Entomology*. 119:55-61.
- Miller MC, Rials TG, Kinn DN, Rials TG. 1994. Chemical stimulation of host tree antibiosis for southern pine beetle, Dendroctonus frontalis, suppression. Symposium on current research in the chemical sciences. Third Annual Southern Station Chemical Sciences Meeting, Alexandria, Louisiana, 7-8 February 1990. General Technical Report Southern Forest Experiment Station, USDA Forest Service. SO-101. Pp 28-32. New Orleans, Louisiana, USA
- Mitchell RJ, Runion GB, Kelley WD, Gjerstad DH, Brewer CH. 1991. Factors associated with loblolly pine mortality on former agricultural sites in the Conservation Reserve Program. *Journal of Soil and Water Conservation*. 46:306-311.
- Moser JC, Bridges JR. 1986. Tarsonemus mites phoretic on the southern pine beetle: attachment sites and numbers of bluestain ascospores carried. *Proceedings of the Entomological Society of Washington*. 88:297-299.
- Nebeker TE, Hodges JD, Blanche CA, Honea CR, Tisdale RA. 1992. Variation in the constitutive defensive system of loblolly pine in relation to bark beetle attack. *Forest Science*. 38:457-466.
- Negron JF, Clarke SR. 1995. Scale insect outbreaks associated with new pyrethroids in a loblolly pine seed orchard. *Journal of Entomological Science*. 30:149-153.
- Nevill RJ, Kelley WD, Hess NJ, Perry TJ. 1995. Pathogenicity to loblolly pines of fungi recovered from trees attacked by southern pine beetles. *Southern Journal of Applied Forestry*. 19:78-83
- Nord JC, Hastings FL, Jones AS. 1990. Field tests of pine oil as a repellent for southern pine bark beetles. Research Note Southeastern Forest Experiment Station, USDA Forest Service. SE-355. 8 pp.
- O' Donnell BP, Payne TL, Walsh KD. 1986. Effect of pine oil on landing and attack by the southern pine beetle. *Journal of Entomological Science*. 21:319-321.
- Oak SW, Tainter FH. 1988. Risk prediction of loblolly pine decline on littleleaf disease sites in South Carolina. *Plant Disease*. 72:289-293.
- Oku H, Shiraishi T, Chikamatsu K. 1989. Active defense as a mechanism of resistance in pine against pine wilt disease. *Annals of the Phytopathological Society of Japan*. 55:603-608.
- Paine TD, Blanche CA, Nebeker TE, Stephen FM. 1987. Composition of loblolly pine resin defenses: comparison of monoterpenes from induced lesion and sapwood resin. *Canadian Journal of Forest Research*. 17:1202-1206
- Paine TD, Stephen FM. 1987. Fungi associated with the southern pine beetle avoidance of induced defense response in loblolly pine. *Oecologia*. 74:377-379
- Paine TD, Stephen FM. 1987. Influence of tree stress and site quality on the induced defense system of loblolly pine. *Canadian Journal of Forest Research*. 17:569-571.
- Paine TD, Stephen FM. 1987. Response of loblolly pine to different inoculum doses of *Ceratocystis minor*, a blue stain fungus associated with *Dendroctonus frontalis*. *Canadian Journal of Botany*. 65:2093-2095.
- Paine TD, Stephen FM. 1987. The relationship of tree height and crown class to the induced plant defenses of loblolly pine. *Canadian Journal of Botany*. 65:2090-2092.
- Paine TD, Stephen FM. 1988. Induced defenses of loblolly pine, Pinus taeda: Potential impact on Dendroctonus frontalis within tree mortality. *Entomologia Experimentalis et Applicata*. 46:39-46.

- Paine TD, Stephen FM, Cates RG. 1988. Phenology of an induced response in loblolly pine following inoculation of fungi associated with the southern pine beetle. *Canadian Journal of Forest Research*. 18:1556-1562.
- Paine TD, Stephen FM, Cates RG. 1993. Within and among tree variation of the induced response of loblolly pine to a fungus associated with Dendroctonus frontalis and sterile wounding. *Canadian Entomologist*. 125:65-71
- Payne TL, Billings RF. 1989. Evaluation of (S I verbenone applications for suppressing southern pine beetle infestations. *Journal of Economic Entomology*. 82:1702-1708
- Peterson GW. 1986. Recent research on conifer needle diseases. General Technical Report, USDA Forest Service, Washington, DC. GTR WO-50. 106pp. Proceedings of conference at Gulfport, MS, Oct. 14-18, 1984.
- Popp MP, Johnson JD, Lesney MS. 1995. Changes in ethylene production and monoterpene concentration in slash pine and loblolly pine following inoculation with bark beetle vectored fungi. *Tree Physiology*. 15:807-812
- Popp MP, Johnson JD, Massey TL. 1991. Stimulation of resin flow in slash and loblolly pine by bark beetle vectored fungi. *Canadian Journal of Forest Research*. 21:1124-1126
- Powers HR Jr.. 1988. Control of tip moth by carbofuran reduces fusiform rust infection on loblolly pine. Research Paper Southeastern Forest Experiment Station, USDA Forest Service. SE-270. 4 pp.
- Powers HR Jr., Waters WE, Ehnstrom B, Ko JH, Kanamitsu K, Lee BY, Sone K, Hong SH, Choi SY, Stone DM, Peng LF, Tsai CY, Chen ZC, La YJ, Yi CK, Saho H, Kim HJ, Yamazaki S, Katsuya K, Sato S. 1985. Abstracts from IUFRO conference on forest gall midges and rusts of pines. Sept. 16-21, 1985, Seoul, Korea. *Journal of Korean Forestry Society*. 71:99-109.
- Powers HR Jr., Belanger RP, Pepper WD, Hastings FL. 1992. Loblolly pine seed sources differ in susceptibility to the southern pine beetle in South Carolina. *Southern Journal of Applied Forestry*. 16:169-174
- Powers HR Jr., Kraus JF. 1986. A comparison of fusiform rust resistant loblolly pine seed sources. *Southern Journal of Applied Forestry*. 10:230-232
- Powers HR Jr., Kuhlman EG. 1987. Resistance of loblolly pine sources to fusiform rust in field progeny tests. Research Paper Southeastern Forest Experiment Station, USDA Forest Service. SE-268. 10 pp.
- Powers HR Jr., Matthews FR. 1987. Five fusiform rust resistant seed sources in coastal South Carolina: a field comparison. *Southern Journal of Applied Forestry*. 11:198-201
- Powers HR, Kraus JF. 1988. Rust resistant loblolly pines. Georgia Forest Research Paper. No.74. 7 pp.
- Powers HR, Miller T, Belanger RP. 1993. Management strategies to reduce losses from fusiform rust. *Southern Journal of Applied Forestry*. 17:146-149
- Pulley PE, Flamm RO, Coulson RN. 1988. Estimating Dendroctonus frontalis daily infestation dynamics. *Researches on Population Ecology*. 30:193-214.
- Riley MA, Goyer RA. 1988. Seasonal abundance of beneficial insects and Ips spp. engraver beetles in felled loblolly and slash pines in Louisiana. *Journal of Entomological Science*. 23:357-365
- Ross DW, Berisford CW. 1990. Nantucket pine tip moth response to water and nutrient status of loblolly pine. *Forest Science*. 36:719-733.
- Ross DW, Berisford CW, Godbee JF Jr.. 1990. Pine tip moth, Rhyacionia spp., response to herbaceous vegetation control in an intensively site prepared loblolly pine plantation. *Forest Science*. 36:1105-1118
- Ross DW, Birgersson G, Espelie KE, Berisford CW. 1995. Monoterpene emissions and cuticular lipids of loblolly and slash pines: Potential bases for oviposition preference of the Nantucket pine tip moth. *Canadian Journal of Botany*. 73:21-25

- Ross DW, Fenn P, Stephen FM. 1992. Growth of southern pine beetle associated fungi in relation to the induced wound response in loblolly pine. *Canadian Journal of Forest Research*. 22:1851-1859
- Rowan SJ. 1987. Nursery seedling quality effects growth and survival in outplantings. *Georgia Forest Research Paper*. No.70. 14 pp.
- Runion GB, Bruck RI. 1988. The effects of thiabendazole on Fusarium subglutinans, the causal agent of pitch canker of loblolly pine. *Plant Disease*. 72:297-300
- Runion GB, Cade SC, Bruck RI. 1993. Effects of carbofuran and thiabendazole on incidence of pitch canker of loblolly pine. *Plant Disease*. 77:166-169
- Runion GB, Kelley WD. 1993. Characterization of a binucleate Rhizoctonia species causing foliar blight of loblolly pine. *Plant Disease*. 77:754-755.
- Salom SM, Grosman DM, McClellan QC, Payne TL. 1995. Effect of an inhibitor based suppression tactic on abundance and distribution of the southern pine beetle and its natural enemies. *Journal of Economic Entomology*. 88:1703-1716.
- Schier GA, Patton RL. 1995. Damping off disease of pine seedlings on soils treated with simulated acidic rain. *Canadian Journal of Forest Research*. 25:838-844.
- Schmidt RA, Allen JE, Belanger RP, Miller T. 1995. Influence of oak control and pine growth on fusiform rust incidence in young slash and loblolly pine plantations. *Southern Journal of Applied Forestry*. 19:151-156
- Schmidt RA, Miller T, Holley RC, Belanger RP, Allen JE. 1988. Relation of site factors in fusiform rust incidence in young slash and loblolly pine plantations in the coastal plain of Florida and Georgia. *Plant Disease*. 72:710-714
- Schmitt JJ, Nebeker TE, Blanche CA, Hodges JD. 1988. Physical properties and monoterpene composition of xylem oleoresin along the bole of Pinus taeda in relation to southern pine beetle attack distribution. *Canadian Journal of Botany*. 66:156-160.
- Shoulders E, Nance WL. 1987. Effects of fusiform rust on survival and structure of Mississippi and Louisiana loblolly pine plantations. *Research Paper Southern Forest Experiment Station, USDA Forest Service*. SO-232. 11 pp.
- Showalter TD, Turchin P. 1993. Southern pine beetle infestation development: interaction between pine and hardwood basal areas. *Forest Science*. 39:201-210.
- Sluder ER. 1993. Results at age 15 years from a half diallel cross among 10 loblolly pines selected for resistance to fusiform rust (Cronartium quercuum f.sp. fusiforme). *Silvae Genetica*. 42:223-230
- Sluder ER, Powers HR Jr.. 1986. Further comparisons between infection of loblolly and slash pines by fusiform rust after artificial inoculation or planting. *Research Note Southeastern Forest Experiment Station, USDA Forest Service*. SE-342. 3 pp.
- Smith AE. 1989. Interference with loblolly pine (Pinus taeda) seedling growth by three grass species. *Weed Technology*. 3:696-698
- Smith MT, Payne TL, Birch MC. 1990. Olfactory based behavioral interactions among five species in the southern pine bark beetle group. *Journal of Chemical Ecology*. 16:3317-3331
- Snow GA, Matthews FR, Nance WL, Foster GS. 1990. Effects of pollen source on loblolly pine resistance to Cronartium quercuum f. sp. fusiforme. *Forest Science*. 1990, 36:304-312
- Solel Z, Bruck RI. 1989. Effect of nitrogen fertilization and growth suppression on pitch canker development on loblolly pine seedlings. *Journal of Phytopathology*. 125:327-335
- Solel Z, Bruck RI. 1990. Relation between wilt rate and obstruction of water flow in stems of two families of loblolly pine affected by pitch canker. *European Journal of Forest Pathology*. 20:317-320

- Solel Z, Runion GB, Bruck RI. 1988. Fusarium equiseti pathogenic to pine. Transactions of the British Mycological Society. 91:536-537
- Spaine P, MacFall JS, Johnson GA, Vozzo JA. 1994. Magnetic resonance microscopy of water movement through fusiform rust galls of pine. Research and applications of chemical sciences in forestry. Proceedings of the 4th Southern Station Chemical Sciences Meeting, Feb. 1-2, 1994, Starksville, MS. General Technical Report Southern Forest Experiment Station, USDA Forest Service. SO-104. Pp.11-16. New Orleans, Louisiana, USA
- Spaine PC. 1987. The development and application of an ELISA for fusiform rust disease resistance screening in vitro in loblolly pine seedlings. Dissertation Abstracts International. B. Sciences and Engineering. 47:4044 B, Thesis, North Carolina State University, USA (1986), 120 pp.
- Stone C, Simpson JA. 1990. Species associations in Ips grandicollis galleries in Pinus taeda. New Zealand Journal of Forestry Science. 20:75-96
- Stone DM, Powers HR. 1989. Sewage sludge increases early growth and decreases fusiform rust infection of nursery run and rust resistant loblolly pine. Southern Journal of Applied Forestry. 13:68-71
- Strom BL, Hain FP. 1996. Host choice of late instar gypsy moths between loblolly pine and sweetgum. Environmental Entomology. 25:603-610
- Strom BL, Hain FP, Ayres MP. 1996. Field performance of F1 sterile gypsy moth larvae on loblolly pine and sweetgum. Environmental Entomology. 25:749-756
- Su KJ, Tan SS, Deng Q. 1991. Studies on symptoms and causative agents of dieback of exotic pines in China. Forest Pest and Disease. 1:2-5.
- Taylor AD, Hayes JL, Roton L, Moser JC. 1992. A phloem sandwich allowing attack and colonization by bark beetles and associates. Journal of Entomological Science. 27:311-316.
- Thompson LC. 1987. Pitch blister moth on loblolly pine. Arkansas Farm Research. 36:5, 10
- Thompson LC, Colvin RJ. 1990. Biological notes on the mealybug Dysmicoccus obesus on loblolly pine in southern Arkansas. Journal of Entomological Science. 25:89-98
- Tisdale RA, Nebeker TE. 1992. Resin flow as a function of height along the bole of loblolly pine. Canadian Journal of Botany. 70:2509-2511
- Twardus D, Taylor S. 1975. Attractant/bait traps for the pales weevil, Hylobius pales: effect of pine species and trap size. Forestry Note Division of Forest Resources, North Carolina. No.22. 15 pp. Raleigh, North Carolina, USA
- Vitousek PM, Andariese SW, Matson PA, Morris L, Sanford RL. 1992. Effects of harvest intensity, site preparation, and herbicide use on soil nitrogen transformations in a young loblolly pine plantation. Forest Ecology and Management. 49:277-292
- Vitousek PM, Matson PA. 1985. Disturbance, nitrogen availability, and nitrogen losses in an intensively managed loblolly pine plantation. Ecology. 66:1360-1376
- von Weissenberg, K. 1990. Host-parasite relationships in forest ecosystems: a review. Silva Fennica. 24:129-139
- Wagner TL, Flamm RO, Hennier PB, Wu W, Coulson RN. 1988. A temperature dependent model of reemergence of Ips avulsus. Environmental Entomology. 17:192-198.
- Walkinshaw CH, Anderson RL. 1988. How to classify fusiform rust galls on infected loblolly and slash pines. Protection Report — Southern Region, USDA Forest Service. R8-PR-14. 4pp.
- Walkinshaw CH, Barnett JP. 1995. Tolerance of loblolly pines to fusiform rust. Southern Journal of Applied Forestry. 19:60-64

- Weatherby JC, Moser JC, Gagne RJ, Wallace HN. 1989. Biology of a pine needle sheath midge, Contarinia acute, on loblolly pine. *Proceedings of the Entomological Society of Washington*. 91:346-349
- Wilcox PL, Amerson HV, Kuhlman KG, Liu BenHui, O' Malley DM, Sederoff RR, Liu BH. 1996. Detection of a major gene for resistance to fusiform rust disease in loblolly pine by genomic mapping. *Proceedings of the National Academy of Sciences of the United States of America*. 93:3859-3864
- Winner WE, Cotter IS, Powers HR Jr., Skelly JM. 1987. Screening loblolly pine seedling responses to SO₂ and O₃: analysis of families differing in resistance to fusiform rust disease. *Environmental Pollution*. 47:205-220
- Yamada, T., and Ito, S. 1993. Chemical defense responses of wilt resistant pine species, Pinus strobus and P. taeda, against Bursaphelenchus xylophilus infection. *Annals of the Phytopathological Society of Japan* 59:666-672.
- Yamada T, Ito S. 1993. Histological observations on the responses of pine species, Pinus strobus and P.taeda, resistant to Bursaphelenchus xylophilus infection. *Annals of the Phytopathological Society of Japan*. 59:659-665
- Yang GY, Li JX, Yang DG, Tan SS. 1988. Investigation on dieback disease of Pinus elliottii and P.taeda in Hunan. *Forest Pest and Disease*. 1:20-22.
- Ye JR, Han ZM, Li CD, Gao SZ. 1986. Resistance to brown spot disease [Mycosphaerella dearnessii] of geographic seed sources of slash pine and loblolly pine [Pinus elliottii, P. taeda] and selection of resistant phenotypes of slash pine individual trees. *Journal of Nanjing Institute of Forestry*. 2: 27-36
- Young SY, Yearian WC. 1987. Intra and inter colony transmission of a nuclear polyhedrosis virus of the loblolly pine sawfly, Neodiprion taedae linearis Ross on pine. *Journal of Entomological Science*. 22:29-34
- Zhu SK. 1988. Control of witches' broom in slash pine and loblolly pine through borax application. *Journal of Zhejiang Forestry Science and Technology*. 8:38-39.
- Zutter BR, Gjerstad DH, Glover GR. 1987. Fusiform rust incidence and severity in loblolly pine plantations following herbaceous weed control. *Forest Science* 33:790-800.
-